




HIV Surveillance – Persons Who Inject Drugs 2018 (preliminary)



Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2010–2017—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises less than 1% of cases.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.





Deaths of Persons with Diagnosed HIV Infection Attributed to Injection Drug Use by Race/Ethnicity, 2017—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	7	0.2
Asian ^a	8	0.2
Black/African American	1,700	47.2
Hispanic/Latino ^b	870	24.1
Native Hawaiian/other Pacific Islander	0	0.0
White	766	21.3
Multiple races	255	7.1
Total^c	3,606	100

Note. Deaths of persons with diagnosed HIV infection may be due to any cause. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.





Persons Living with Diagnosed HIV Infection Attributed to Injection Drug Use by Race/Ethnicity, Year-end 2017—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	481	0.4
Asian ^a	611	0.5
Black/African American	57,641	46.6
Hispanic/Latino ^b	33,466	27.0
Native Hawaiian/other Pacific Islander	46	0.0
White	25,770	20.8
Multiple races	5,629	4.5
Total^c	123,798	100

Note. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Includes Asian/Pacific Islander legacy cases.


^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total. Includes 155 persons whose race/ethnicity is unknown.





Diagnosed HIV Infections Classified as Stage 3 (AIDS) among Persons Who Inject Drugs, 1985–2017—United States and 6 Dependent Areas




Note. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.





Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Transmission Category and Year of Classification, 1985–2017

United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.





Diagnosed HIV Infections Classified as Stage 3 (AIDS) among Persons Who Inject Drugs, by Race/Ethnicity and Year of Classification, 1985–2017—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.





Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS) and Who Inject Drugs, Year-end 2017—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	254	0.3
Asian ^a	328	0.4
Black/African American	37,030	47.4
Hispanic/Latino ^b	22,458	28.5
Native Hawaiian/other Pacific Islander	24	0.0
White	14,698	18.6
Multiple races	4,020	5.1
Total^c	78,816	100

Note. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Includes Asian/Pacific Islander legacy cases.


^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total. Includes 5 persons whose race/ethnicity is unknown.





Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Transmission Category, 2018—United States and 6 Dependent Areas



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. "Other" transmission category not displayed as it comprises less than 1% of cases.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



Diagnoses of HIV Infection, by Age Group, Race/Ethnicity, and Injection Drug Use-Related Transmission Category, 2018—United States and 6 Dependent Areas

Race/ethnicity	Age group (years), %					Total
	13–19 N=72	20–24 N=370	25–34 N=1,341	35–44 N=912	≥45 N=1,132	
American Indian/Alaska Native	2.1	0.6	1.5	0.7	1.0	1.1
Asian	0.0	0.7	1.0	0.8	0.7	0.8
Black/African American	33.2	25.3	21.4	23.2	36.3	26.9
Hispanic/Latino ^a	23.0	28.6	21.1	19.9	21.7	21.8
Native Hawaiian/other Pacific Islander	0.0	0.5	0.2	0.0	0.2	0.2
White	33.8	40.4	51.0	52.9	37.3	46.0
Multiple races	7.9	3.8	3.9	2.6	2.8	3.3
Total^b	100	100	100	100	100	100

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Two standard transmission categories are collectively referred to as related to injection drug use (IDU-related), and these two categories are mutually exclusive: injection drug use; and male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.

^b Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.





Diagnoses of HIV Infection, by Age Group, Sex, and IDU-Related Transmission Category, 2018—United States and 6 Dependent Areas


Transmission category	Age group (years), %					Total
	13–19 N=72	20–24 N=370	25–34 N=1,341	35–44 N=912	≥45 N=1,132	
Male injection drug use	21.8	24.7	32.5	40.6	48.5	38.2
Male-to-male sexual contact and injection drug use	46.5	48.6	44.4	29.6	21.0	34.4
Female injection drug use	31.7	26.7	23.2	29.8	30.5	27.4
Total^a	100	100	100	100	100	100

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

^a Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Persons Who Inject Drugs, by Sex and Race/Ethnicity, 2018—United States and 6 Dependent Areas



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

^aHispanics/Latinos can be of any race.





Diagnoses of HIV Infection Among Persons Who Inject Drugs, by Race/Ethnicity 2018—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	21	0.8
Asian	18	0.7
Black/African American	770	30.7
Hispanic/Latino ^a	511	20.4
Native Hawaiian/other Pacific Islander	2	0.1
White	1,115	44.4
Multiple races	76	3.0
Total^b	2,511	100

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.

^b Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.





Diagnoses of HIV Infection Among Males Who Inject Drugs, by Race/Ethnicity 2018—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	7	0.5
Asian	12	0.8
Black/African American	450	30.8
Hispanic/Latino ^a	362	24.8
Native Hawaiian/other Pacific Islander	2	0.1
White	593	40.6
Multiple races	37	2.5
Total^b	1,462	100

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.

^b Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.





Diagnoses of HIV Infection Among Females Who Inject Drugs, by Race/Ethnicity 2018—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	14	1.3
Asian	6	0.6
Black/African American	320	30.5
Hispanic/Latino ^a	149	14.2
Native Hawaiian/other Pacific Islander	0	0.0
White	522	49.8
Multiple races	39	3.7
Total^b	1,049	100

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.


^a Hispanics/Latinos can be of any race.

^b Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.





HIV Diagnoses Among Persons Who Inject Drugs, by Region and Race/Ethnicity 2018—United States and 6 Dependent Areas




Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.





Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Sex and Transmission Category, 2018—United States and 6 Dependent Areas



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.





Diagnosed HIV Infections Classified as Stage 3 (AIDS) among Males Who Inject Drugs, by Race/Ethnicity, 2018—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	7	0.8
Asian ^a	7	0.8
Black/African American	380	41.1
Hispanic/Latino ^b	249	26.9
Native Hawaiian/other Pacific Islander	0	0.0
White	243	26.3
Multiple races	37	4.0
Total^c	925	100

Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.





Diagnosed HIV Infections Classified as Stage 3 (AIDS) among Females Who Inject Drugs, by Race/Ethnicity, 2018—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	5	0.7
Asian ^a	4	0.5
Black/African American	298	40.5
Hispanic/Latino ^b	126	17.1
Native Hawaiian/other Pacific Islander	0	0.0
White	255	34.6
Multiple races	47	6.4
Total^c	736	100

Note. Data for the year 2018 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

^a Includes Asian/Pacific Islander legacy cases.


^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.





Diagnosed HIV Infections Classified as Stage 3 (AIDS) among Persons Who Inject Drugs, by Region and Race/Ethnicity, 2018—United States and 6 Dependent Areas



Note. Data for the year 2018 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data exclude men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

